

Spectrophotometric determination of total alkaloids in some Iranian medicinal plants

Shamsa Fazel¹, Monsef Hamidreza², Ghamooshi Rouhollah² and Mohammadreza Verdian-rizi^{2*}

¹Department of Medicinal Chemistry, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran.

²Department of Pharmacognosy, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran.

*E-mail: mverdian@razi.tums.ac.ir

Abstract

A simple spectrophotometric method based on the reaction with Bromocresol Green (BCG) was developed for determination of total alkaloids in medicinal plants. A yellow complex forms and is easily extractable by chloroform at pH 4.7. The absorbance of the complex obeys Beer's law over the concentration range of 4-13 µg atropine per mL of chloroform. This procedure can be carried out in the presence of other compounds without interference.

Key words: BCG, total alkaloids, medicinal plants, determination

Introduction

The alkaloids represent a group of natural products that had a major impact on the economic, medical and social affairs of humans. Many of these agents have potent physiological effects on mammalian systems as well as other organisms, and as a consequence, some constitute important therapeutic agents. Atropine, morphine, quinine and vincristine are representative of a host of agents used to treat a range of disease conditions that range from malaria to cancer. Therefore determination of total alkaloids is very important related to the quality of medicinal plants (Robbers *et al.*, 1996).

The methods reported for the determination of alkaloids include official methods (Kartal, 2001; Levent, 2002;), high-performance liquid chromatography (HPLC) (Tomonari *et al.*, 1994; Salvadori *et al.*, 1994; Qi *et al.*, 2002; Kanazawa *et al.*, 2000), fluorimetry (Masatoki *et al.*, 1989; Andio *et al.*, 1987), ion-chromatography (Qing-Chun *et al.*, 2001), coulometry (Qing-qin *et al.*, 2002), gas-chromatography (Pagliariussi *et al.*, 2002; Chernyseva *et al.*, 2001), and electrochromatography. Most of the reported spectrophotometric methods have disadvantages such as narrow range of determination, require heating or extraction, a long time is needed for the reaction to be completed, and the coloured product formed is unstable. The purpose of the current work was to provide a simple, sensitive, and rapid spectrophotometric method for the determination of total alkaloids in medicinal plants. The method is based on the reaction of alkaloid with Bromocresol Green, forming a yellow-colored product. The method offers the advantages of sensitivity and stability.

Materials and methods

Plant material: Plant material including *Acroptilon repens* L. (aerial parts), *Berberis vulgaris* L. (aerial parts, fruits), *Biebersteinia multifida* DC. (aerial parts, root), *Calendula officinalis* (flower), *Chelidonium majus* L. (aerial parts), *Echium amoenum* Fish & Mey (flower), *Equisetum arvense* L. (aerial

parts), *Hyoscyamus niger* L. (aerial parts), *Hypocoum pendulum* L. (aerial parts), *Malva sylvestris* L. (aerial parts), *Scrophularia striata* Bioss. (root) and *Stachys lavandulifolia* Vahl. (aerial parts), was collected from local market of Tehran province, in May 2003. All plants were identified in the Herbarium of Faculty of Pharmacy, Tehran University of Medical Sciences.

Bromocresol green solution (1×10⁻⁴): Bromocresol green (69.8 mg) was warmed with 3 mL of 2N NaOH and 5 mL distilled water until completely dissolved and diluted to 1000 mL with distilled water.

Phosphate buffer (pH 4.7): pH of 2 M sodium phosphate (71.6 g Na₂HPO₄ in 1 L distilled water) was adjusted to pH 4.7 with 0.2 M citric acid (42.02 g citric acid in 1 L distilled water).

Atropine standard solution: One mg pure atropine (Sigma Chemical Co.) was dissolved in 10 mL distilled water.

Preparation of standard curve: Accurately measured aliquots (0.4, 0.6, 0.8, 1 and 1.2 mL) of atropine standard solution were transferred to different separatory funnels. Five mL phosphate buffer (pH 4.7) and 5 mL BCG solution were added. Mixture was shaken with 1, 2, 3 and 4 mL of chloroform. The extracts were collected in a 10 mL volumetric flask and then diluted to volume with chloroform. The absorbance of the complex in chloroform was measured at 470 nm against blank prepared as above but without atropine.

Extraction: The plant material (100g) was grinded and then extracted with methanol for 24 h in a continuous extraction (soxhlet) apparatus. The extract was filtered and methanol was evaporated on a rotary evaporator under vacuum at a temperature of 45°C to dryness. A part of this residue was dissolved in 2 N HCl and then filtered. One mL of this solution was transferred to a separatory funnel and washed with 10 mL chloroform (3 times). The pH of this solution was adjusted to neutral with 0.1 N NaOH. Then 5 mL BCG solution and 5 mL phosphate buffer were added to this solution. The mixture was shaken and the complex formed

Specimen Copy: Not for sale

was extracted with 1, 2, 3, and 4 mL chloroform by vigorous shaking. The extracts were collected in a 10 mL volumetric flask and diluted to volume with chloroform. The absorbance of the complex in chloroform was measured at 417 nm vs. similarly prepared blank.

Results and discussion

A yellow-coloured complex with a maximum absorption was developed. This complex was completely extractable by chloroform at pH 4.7. A calibration curve was plotted for various concentration of atropine (Fig.1). Beer's law was followed over the concentration range of 4-13 µg atropine per mL of chloroform. The effect of temperature and pH were studied. A pH of 4.7 gave optimum results and different temperatures had no effect on complex formation and extraction. The complex was very stable in chloroform and began to fade slowly only after 10 days. Before the extraction, the mixture was put in a boiling water bath for 3 min. The absorbance did not change after extraction with chloroform. Table 1 shows the amount of total alkaloid in tested plant materials determined by BCG-complex formation method.

Table 1. Determination of total alkaloids in tested plant materials (100g) by BCG-complex formation

No.	Plant	Part used	Amount (mg)	Amount (M mol)
1	<i>Acroptilon repens</i> L.	Aerial parts	13.35	0.023
2	<i>Berberis vulgaris</i> L.	Aerial parts	40.58	0.070
3	<i>Berberis vulgaris</i> L.	Fruit	19.70	0.034
4	<i>Biebersteinia multifida</i> DC.	Aerial parts	204.56	0.353
5	<i>Biebersteinia multifida</i> DC.	Root	1688.47	2.920
6	<i>Calendula officinalis</i> L.	Flower	16.14	0.028
7	<i>Chelidonium majus</i> L.	Aerial parts	248.09	0.430
8	<i>Echium amoenum</i> Fish & Mey	Flower	18.44	0.320
9	<i>Equisetum arvense</i> L.	Aerial parts	255.02	0.440
10	<i>Hyoscyamus niger</i> L.	Aerial parts	324.09	0.560
11	<i>Hypocoum pendulum</i> L.	Aerial parts	39.20	0.068
12	<i>Malva sylvestris</i> L.	Aerial parts	35.06	0.060
13	<i>Scrophularia striata</i> Bioss.	Root	7.90	0.014
14	<i>Stachys lavandulifolia</i> Vahl.	Aerial parts	9.73	0.017

A few methods with different sensitivities have been developed for determination of alkaloids in plant materials for example gravimetric and titrimetric methods. These methods lack the adequate sensitivity and have some problems. As with most gravimetric methods, the residue obtained is found to be impure since more than one spot is revealed by TLC. In titrimetric assay, the end-point is masked by the color of the extract. On the other hand, there is no constant method applicable for all alkaloids. Methods with high sensitivity such as HPLC are not routine methods for determination of total alkaloids and these methods are very costly and need special equipment. Spectrophotometric determination of total alkaloids with bromocresol green is a simple and sensitive method and do not need very special equipment. The proposed method has the advantage of being less time consuming, with the assay requiring an average of 1 h.

Acknowledgement

We are grateful to the Faculty of Pharmacy, Tehran University of Medical Sciences, for the financial support of this investigation.




Fig 1. Variation of the absorbance vs. atropine concentration at 470 nm

References

- Andio, M.M., C.G. De Lima and J.D. Winefordner, 1987. Luminescence characteristics of caffeine and theophylline. *Spectrochim. Acta A*, 43: 427-430.
- Chernyseva, N.N., I.F. Abdullin and G.K. Bundikov, 2001. Coulometric determination of purine alkaloids series with electrogenerated chlorine. *J. Anal. Chem.*, 56: 663-665.
- Kanazawa, H., J. Kizu and Y. Matsushima, 2000. Simultaneous determination of theophylline and its metabolites by HPLC. *Yakagaku Zasshi*, 120: 1051-1060.
- Kartal, M. 2001. LC method for the analysis of paracetamol, caffeine, and codeine phosphate in pharmaceutical preparations. *J. Pharm. Biomed. Anal.*, 26: 857-864.
- Levent, A.M. 2002. HPLC method for the analysis of paracetamol, caffeine, and dipyrone. *Turk. J. Chem.*, 26: 521-528.
- Masatoki, K. and T. Hirokazu, 1989. Fluorometric reactions of purines and determination of caffeine. *Talanta*, 36: 1171-1175.
- Pagliariussi, R., S. Frietas, A.P. Luis and J.K. Bastos, 2002. A quantitative method for the analysis of xanthine alkaloids in *Paullinia cupana* (guarana) by capillary gas chromatography. *J. Sep. Sci.*, 25: 371-374.
- Qi, M.L., P. Wang, Y.X. Lang, J.L. Lang and R.N. Fu, 2002. Simultaneous determination of caffeine, theophylline, and theobromine by HPLC. *J. Chromatogr. Sci.*, 40: 45-48.
- Qing-Chun, C. and J. Wang, 2001. Simultaneous determination of artificial sweeteners, preservatives, caffeine, theobromine, and theophylline in food and pharmaceutical preparations by ion chromatography. *J. Chromatogr. A*, 937: 57-64.
- Qing-qin, X., D.L. Ming, J.P. Wang and A.H. Bai, 2002. Direct determination of caffeine and theophylline by gas chromatography. *Fenxi Kexue Xuebao*, 18: 520-525.
- Robbers, E., K.M. Speedie, V.E. Tyler, 1996. *Pharmacognosy and Phramacobiotechnology*, Baltimore : Williams and Wilkins, 40-45.
- Salvadori, M.C., E.M. Reiser and L.M. Ribeironeta, 1994. Determination of xanthines by HPLC and TLC in horse urine after ingestion of guarana powder. *Analyst*, 119: 2701-2703.
- Tomonari, U., K. Ryuii, T. Kazami and H. Hiraki, 1994. Direct injection determination of theophylline and caffeine in blood serum by HPLC using an ODS column coated with zwitterionic bile acid derivative. *Analyst*, 119: 1767-1770.