

# The role of natural exogenous foliar applications in alleviating salinity stress in *Lagerstroemia indica* L. seedlings

Amira Sh. Soliman<sup>1\*</sup> and Nermeen T. Shanan<sup>2</sup>

<sup>1</sup>Natural Resources Department, Institute of African Research and Studies, Cairo University, Giza, Egypt. <sup>2</sup>Ornamental Horticulture Department, Faculty of Agriculture, Cairo University, Giza, Egypt. \*E-mail: sitamira2000@yahoo.com.

## Abstract

A pot experiment, conducted during 2014 and 2015 seasons in completely randomized factorial design to determine the effect of natural extracts, foliar spray of *Moringa* leaves extract (1:30), humic acid (10%), seaweed (2%), Hogland nutrient solution and tap water as control, on growth characteristics (plant height, stem diameter, number of leaves/plant, number of branches/plant, root length, and total dry weight of plant parts (roots, shoots and flowers), floral and chemical characteristics of *Lagerstroemia indica* grown at various sea salt concentrations (0, 3.12, 6.25 and, 9.37 dS/m) showed that by increasing sea salt concentrations, all growth characteristics, inflorescence number/plant and, inflorescence diameter decreased significantly, while, the number of days to inflorescence increased. Total chlorophylls, carotenoid contents, total carbohydrates and N, P, K%. Meanwhile, proline content, total soluble phenols, Na, Ca, and antioxidant enzymes (catalase, superoxide dismutase and peroxidase) increased at the high level of salinity (9.37 dS/m). The usage of the *Moringa* leaf extract significantly improved growth, inflorescence, as well as chemical characteristics, but also, decreased significantly Na under the adverse conditions of the studied sea salt stress. *Moringa* leaf extract could promote growth and protect crape myrtle plants against injuries by sea salt stress and being safe and cheap can substitute inorganic or chemical fertilizer.

**Key words:** *Lagerstroemia indica*, foliar applications, *Moringa* leaves extract, sea salt stress, inflorescence characteristics, antioxidant enzymes.

## Introduction

*Lagerstroemia indica* L. (Crape myrtle) is one of the most famous species of the *Lagerstroemia* genus, which belongs to the *Lythraceae* family (Conti *et al.*, 1997) and includes 53 woody species (Tobe and Raven, 1990). It is a widely commercialized as a deciduous shrub or small tree (Anonymous, 1962) that has become naturalized and invasive in many tropical and subtropical regions of the world (Randall, 2012).

Crape myrtle is utilized as a versatile landscaping plant, decorative, an ornamental and attractive shrub in the world (Anderson, 2007). *L. indica* is used often around homes and in small parking as shade trees, also used for buffer strips along highways, decks, patios, and around parking (Moore and Walker, 2014). Its wood is hard and useful as a timber (Gamble, 1972). *L. indica* has a very dense and wide-spreading root system, so it is used in erosion control. It is also grown as a boundary or barrier support plant in gardens and cultivated areas (Orwa *et al.*, 2009).

It is also a medicinal shrub or small tree (Chopra *et al.*, 1956). The leaves, flowers, roots and bark of *L. indica* are used as purgative, hydrogogue, febrifuge, stimulant and styptic. The leaves have great medicinal value, particularly in the treatment of diabetes. The roots are astringent and used as gargle and seeds contain a narcotic principle and are used in the production of pesticides. This plant is secondary food plants of non-mulberry Tasar silk worm. A paste of the flowers is applied externally to cuts and wounds, while, a decoction of the flowers is used in the treatment of colds (Yeung, 2010; Masharani and German, 2011). Yang *et al.* (2011) found that *L. indica* may be used as a valuable agent for treating allergic disease such as asthma due

to its anti-inflammatory property. Diab *et al.* (2012) found that the methanol extracts of *L. indica* leaves exhibited antimicrobial activity against all pathogenic bacteria and yeast. In addition, Kotnala *et al.* (2013) concluded that particular silent disease can be treated by using extracts from different *Lagerstroemia* species.

*L. indica* is effectively grown well in harsh urban conditions. It is tolerant to fire, drought, cold conditions, and relatively free of disease and insect and thus has a potential to grow in new habitats, displacing and smothering native vegetation and makes it an excellent landscape plant (Foxcroft *et al.*, 2007; Oviedo *et al.*, 2012). However, the literature survey conducted to identify biotic stress (*i.e.* salinity) causing injury and affecting its nursery and landscape industries has indicated crape myrtle as a salt-sensitive species (till 3.0 dS/m) (Cabrera and Devereaux, 1999). Salinity is more apparent in semi-arid and arid regions of the world and responsible for decline in plant growth by osmotic stress, and specific ion toxicity. Therefore, interventions can be made to minimize the effect of salt stress on plants, especially during the early development, to promote successful establishment and growth.

Foliar application with the exogenous plant growth regulators (PGRs) such as cytokinin or antioxidants (ascorbic acid) is the most effective approach involved in promoting plant growth and development by minimizing or alleviating the adverse effects of salt on plant growth and metabolism in many plants (Sadak and Dawood, 2014), but this application is expensive. Therefore, there is a need to explore the natural sources of PGRs for exogenous use which are not only organic, but also easily adapted, environmental friendly, and inexpensive such as extract obtained from *Moringa* (*Moringa oleifera*) leaves (Moussa and Hassan, 2016), humic acid

(HA) (Rady *et al.*, 2016), seaweed extract (SE) (Battacharyya *et al.*, 2015).

Therefore, this investigation was undertaken to evaluate the effects of foliar application of *Moringa* leaves extract, humic acid or seaweed extract to negate the negative effect of salinity on growth, floral and chemical characters of *L. indica*.

## Materials and methods

The pot experiment was carried out at the Experimental Laboratory in the Natural Resources Department, Institute of African Research and Studies, and at the Ornamental Horticulture Department, Faculty of Agriculture, Cairo University, Giza, Egypt, during two successive seasons of 2014 to 2015. The soil texture of pot media was sandy with the following characteristics: coarse sand 30.82%, fine sand 62.61, silt 1.22%, clay 5.35%, pH 7.75, EC 1.15 dS/m, organic matter 0.08%, available N 6.9 ppm, available P 6.2 ppm, available K 64 ppm, CaCO<sub>3</sub> 0.26%, and water holding capacity 14.5%. The soil analysis was carried out according to the methods described by Chapman and Pratt (1961).

**Plant material:** Three years old seedlings of *L. indica* (Dark purple flower color) were obtained from a private nursery in El-Menoufia Governorate, Cairo, Egypt, on the 1<sup>st</sup> of April, in both seasons. The seedlings (15 cm tall) were transplanted into 25 cm diameter-plastic bags filled with 6 kg of the same sandy soil, and watered every 3 days. Recommended dose of chemical fertilizers were added at the rate of 5 g per pot ammonium nitrate (33.5% N), calcium superphosphate (15.5% P<sub>2</sub>O<sub>5</sub>) and potassium sulphate (48-50% K<sub>2</sub>O) for plant maintenance.

**Moringa leaves extract (MLE):** The fresh young *Moringa oleifera* leaves were air dried, ground and extracted according to Price (2007). Supernatant was diluted 30 times with water and used as a foliar spray (Table 1).

**Seaweed extract (SW):** Seaweed extract (powerful) was obtained from Cairochem for Agricultural Services Company, Alexandria, Egypt. The extract contains minerals, vitamins, enzymes, amino acids, sugars, and plant hormones (*i.e.*, auxins, cytokinins and gibberellins) (Table 2).

**Humic acid extract (HA):** Natural humic substances (Table 3) obtained from Soil, Water and Environment Research Institute, Agriculture Research Center (A.R.C).

**Hoagland nutrient solution (HNS):** Full strength Hoagland nutrient solution was prepared with tap water (Hoagland and Arnon, 1950), and the pH of the solution was monitored several times and maintained within a range of 6.0-6.5, which obtained from El-Gomhorya Company, Egypt (Table, 4).

**Treatments:** After 15 days of transplanting, the seedlings were subjected to following different levels of saline water irrigation: non saline water (the control; 0.5 dS/m), saline water 3.12, 6.25 and 9.37 dS/m. Sea salt solutions for irrigation were prepared by adding the required amount of sea salt to tap water per liter, which is a mixture of synthetic seawater salt obtained from Sigma Company. Seedlings were irrigated with gradual increasing sea salt concentration weekly up to reaching the desired salinity levels of the experiment mentioned above. To maintain the required soil medium salt level, the EC of the soil medium was measured

Table 1. Chemical composition of *M. oleifera* leaves extract

Constituent	Content mg g <sup>-1</sup> DW
K	29.00
P	6.15
Ca	9.02
Mg	6.04
Fe	1.90
Zn	0.49
Soluble phenols	2.13
Ascorbic acid	3.48
Amino acids	125.20
Proline	26.19
Total soluble sugars	157.80
Phytohormones	Content (µg g <sup>-1</sup> DW)
Indole-3-acetic acid	0.91
Gibberellins	0.81
Zeatin	0.95
Abscisic acid	0.25

Table 2. Chemical composition of seaweed extract (powerful)

Constituent	Content (%)
<i>Ascophyllum nodosum</i>	75.00
N	1.50
P <sub>2</sub> O <sub>5</sub>	1.00
K <sub>2</sub> O	1.50
S	1.60
Mg	1.60
Ca	1.60
Se	0.02
Mannitol	2
Topolin (6-[3-hydroxy benzyl-amino] purine)	0.01
O.M.	25
Free amino acid	10
Alginate acid	9

periodically by portable EC meter.

Seedlings grown under normal or saline conditions were sprayed four times at 30 day intervals at a rate of 5 mL/ pot, starting on 15<sup>th</sup> May, in both seasons, with each of the following: MLE (1:30), seaweed (2%), HA (10 %), full strength Hoagland nutrient solution as a chemical fertilization and distilled water only as a control. The spraying was done manually using a spraying bottle, on both sides of the leaves evenly and, it was carried out between 09:00 and 11:00 AM. The concentrations of MLE, HA and seaweed sprays were based on results from a preliminary pot trial (data not shown). After 125 days of treatments, the plants were harvested, in both seasons, measured and analyzed. The experiment was arranged as factorial including Randomized Complete Block Design with 20 treatments which were the combinations of four salinity levels (0, 3.12, 6.25 and 9.37 dS/m)

Table 3. Composition of natural humic acid

Chemical	Content (%)
Total Nitrogen (N)	1.40
Phosphorus (P)	0.76
Potassium (K)	2.80
Calcium (Ca)	1.00
Magnesium (Mg)	0.50
Sulfur (S)	1.30
Iron (Fe)	1.39
Boron (B)	0.07

Table 4. Composition of Hoagland nutrient solution

Chemical	Concentration
CaNO <sub>3</sub>	4.0 mM
MgSO <sub>4</sub>	2.0 mM
KNO <sub>3</sub>	4.0 mM
(NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub>	0.4 mM
MnSO <sub>4</sub>	2.0 µM
CuSO <sub>4</sub>	0.3 µM
ZnSO <sub>4</sub>	0.8 µM
NaCl	30.0 µM
Na <sub>2</sub> Mo <sub>4</sub>	0.1 µM
KH <sub>2</sub> PO <sub>4</sub>	1.43 µM
H <sub>3</sub> BO <sub>3</sub>	10.0 µM
Fe-Na-EDTA	20.0 µM

and five foliar applications (MLE, seaweed, HA, Hoagland nutrient solution and the control). Each treatment was replicated six times with one seedling per unit.

**Growth characteristics:** After 60 days from sowing, plants from each replicate were harvested, and the following characters were recorded; plant height (cm), stem diameter (cm), number of leaves/plant, number of branches/plant, root length (cm), and total dry weight of the whole plant (roots, shoots and flowers) (g).

**Floral characteristics:** Number of days to flowering (day), inflorescence number /plant and, inflorescence diameter (cm) were recorded.

#### Chemical analysis

**Photosynthetic pigments and total carbohydrates:** Total chlorophylls and carotenoid contents (mg/g fresh weight) were measured by Spectrophotometer and calculated according to equation described by Moran (1982). The total carbohydrate percentage was determined according to the method of DuBois *et al.* (1956).

**Proline content and total soluble phenols (TSP):** The proline concentration was determined using fresh material according to Bates *et al.* (1973). In ethanol extract, total soluble phenols (mg/g f.w.) was estimated using the Folin-Ciocalteau calorimetric method (Swain and Hillis, 1959).

**Macro-nutrient:** Dried leaf samples were digested as described by Piper (1950) and the extract was analyzed to determine: nitrogen % by the modified micro Kjeldahl method as described

by Pregl (1945), phosphorus % as in Jackson (1967), potassium and sodium were determined according to the method described by (AL-Khayri, 2002) using a flame spectrophotometer, and Ca % was determined by atomic absorption (Allen *et al.*, 1984).

**Antioxidant enzyme determination:** Antioxidant enzyme extraction were carried out using fresh leaf tissues at 40 °C in a buffer solution (3: 1 buffer: fresh weight v/v) in a pastel. It was mortared with 100 mM potassium phosphate buffer (at pH 7.5) containing 1 mM EDTA, 3 mM DL-dithiothreitol and 5% (w/v) insoluble polyvinyl pyrrolidone. The homogenates were centrifuged at 10000 g for 30 min and then the supernatants were stored in separate aliquots at 8 °C (Vitoria *et al.*, 2001). Antioxidant enzymes were assayed as follows; catalase (CAT) by measuring the decrease in absorbance due to disappearance of H<sub>2</sub>O<sub>2</sub> at 240 nm according to Chance and Maely (1955), superoxide dismutase (SOD) by photochemical method as described by Giannopolitis and Ries (1977) and, Ascorbate peroxidase (APX; EC 1.11.1.11) activity was determined as suggested by Rao *et al.* (1996) and the optical density at 290 nm. Enzymes activities were expressed as units/min/mg protein.

**Statistical analysis:** The obtained results were subjected to statistical analysis of variance and the means were compared using the least significant difference (L.S.D.) test at the 5% level, as described by Little and Hills (1978).

## Results and discussions

**Growth characteristics:** Growth characters [plant height, stem diameter, number of leaves/ plant, number of branches/ plant, root length, and total dry weight of the whole plant] of *L. indica* under different sea salt concentrations without any foliar application were reduced (Table 5). Total dry biomass of crape myrtle plants treated with sea salt solutions of 6.25 dS/m and 9.37 dS/m decreased to 27.93 and 23.14 g/plant, in both seasons, respectively, in comparison with non-saline (37.07 g/plant) in both seasons.

The reduction in growth characteristics under sea salt stress conditions may be due to ion toxicity and low water potential which may cause one or more of the following; water deficit, imbalance in absorption, uptake and transport of essential nutrients, turgor pressure reduction in expanding tissues, reduction in photosystem activity in leaf cells, reduced ability to produce and utilize assimilates to the growing regions, and direct effects of accumulated salts on metabolic steps in dividing and expanding cells (Tester and Davenport, 2003; Munns *et al.*, 2006).

A significant increase in growth characteristics was observed in plants sprayed with full strength Hoagland nutrient solution (HNS), seaweed (SW, 2%) or humic acid (HA, 10%), while the foliar spray of *Moringa* leaves extract (MLE) had significant effect, irrespective of growth characteristics under non saline (control) or different sea salt concentrations, which the sea salt concentration at 9.37 dS/m recorded the highest total dry weight (23.30 g/plant) compared to control (10.67g/plant) in the 1<sup>st</sup> season, while in the 2<sup>nd</sup> season (25 g/plant) compared to the control (12.74 g/plant). The effect of foliar application by MLE to improve the plant tolerance to abiotic stresses, including salinity has been reported by many researchers (Howladar, 2014; Rady and Mohamed, 2015).

Table 5. Growth characteristics of *L. indica* affected by exogenous foliar applications under different non-saline and sea salt concentrations during two seasons, 2014 (1<sup>st</sup>) and 2015 (2<sup>nd</sup>).

Treatments	Plant height (cm)		Stem diameter (cm)		Number of leaves/plant		Number of branches/plant		Root length (cm)		Total dry weight (g)		
	1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>	
S0	C	70.33	72.00	1.07	1.30	17.33	19.67	5.00	7.33	46.67	50.00	17.07	20.00
	HNS	84.33	85.67	1.77	1.93	27.67	28.33	11.67	14.33	56.67	59.67	28.03	28.73
	MLE	91.00	96.33	2.77	3.03	31.33	33.00	17.33	18.33	65.00	70.00	31.27	33.13
	SE	84.67	86.67	1.93	2.13	29.67	30.67	14.67	16.33	61.67	65.67	28.53	29.63
	HA	83.33	84.33	1.73	1.87	27.33	28.00	10.33	11.67	55.00	57.33	28.07	28.07
S1	C	68.33	69.67	0.97	1.23	15.67	18.33	4.33	6.00	44.00	48.33	16.23	18.63
	HNS	83.00	84.00	1.60	1.73	24.67	25.33	9.00	9.67	54.33	58.33	24.23	24.70
	MLE	90.00	95.00	2.53	2.73	29.33	31.67	16.00	17.33	62.33	67.67	29.70	31.13
	SE	83.33	85.33	1.70	2.07	27.00	29.67	14.00	14.33	58.33	63.00	27.53	29.27
	HA	81.67	82.33	1.57	1.87	25.00	26.33	9.33	10.33	54.00	59.00	25.47	26.23
S2	C	63.33	66.67	0.83	1.03	12.67	16.00	3.67	5.00	38.00	41.67	12.50	15.43
	HNS	77.67	80.00	1.07	1.20	16.33	22.33	8.00	8.67	47.33	51.67	21.23	22.60
	MLE	84.67	86.33	1.63	1.73	23.67	28.00	14.00	16.00	56.00	60.00	26.27	27.23
	SE	81.00	83.00	1.43	1.47	20.67	26.33	11.33	12.67	52.00	57.00	24.67	24.77
	HA	79.67	81.00	1.13	1.30	19.33	24.33	9.00	9.33	50.00	52.67	22.17	23.83
S3	C	51.67	58.00	0.57	0.83	7.67	11.67	3.00	4.67	32.67	36.67	10.67	12.47
	HNS	61.33	62.67	0.87	1.00	11.67	15.00	7.00	7.67	41.67	43.00	17.10	17.77
	MLE	77.33	79.67	1.30	1.47	17.00	22.67	11.67	12.67	51.00	54.00	23.30	25.00
	SE	74.67	75.67	1.10	1.17	14.67	19.67	10.67	11.00	47.67	49.33	21.40	21.97
	HA	71.67	73.33	0.97	1.07	12.33	16.67	8.00	8.67	43.00	45.00	17.97	18.77
LSD ( $P=0.05$ )		2.74	3.61	0.20	0.23	1.40	1.92	1.08	1.01	3.84	3.08	1.36	1.67
		2.06	1.60	0.15	0.14	1.73	1.85	1.04	0.96	2.49	2.82	1.11	1.24
		4.12	3.19	0.31	0.29	3.47	3.71	2.09	1.93	4.99	5.64	2.23	2.49

S0=control, S1= 3.12 dS/m, S2= 6.25 dS/m, S3= 9.37 dS/m.

C= distilled water as control, HNS= Hoagland nutrient solution, MLE= *Moringa* leaves extract, SE= seaweed, HA= humic acid.

The reason for growth acceleration of *L. indica* plants sprayed by *Moringa* leaves extract (MLE) under normal or saline conditions might be due to the presence of essential macro- and micro-nutrients such as Ca, K, and Zn, and enriched content of phytohormones (indole-3-acetic acid, gibberellins and zeatin as a cytokinin) in MLE that favored rapid division, multiplication and enlargement of plant cells, which encouraged plant growth and productivity under salt stress conditions. It is also rich in crude proteins which are essential for the protoplasm formation (Moyo *et al.*, 2011; Afzal *et al.*, 2012). In addition, Azra *et al.* (2012) and Rehman *et al.* (2014) concluded that MLE applications maintained optimum tissue water status and membranes stabilities, enhanced antioxidant levels and activated plant defense system, increased levels of plant secondary metabolites, reduced uptake of undesirable  $\text{Na}^+$  and/or  $\text{Cl}^-$ , and enhanced leaf  $\text{K}^+$ . All these events lead to vigorous seedling growth, maximizing the crop performance.

**Floral characteristics:** Salt stress significantly affected the floral characteristics (number of days to flowering, inflorescence number /plant and, inflorescence diameter) of *L. indica* plants (Fig. 1). Increasing sea salt concentrations to 9.37 dS/m increased the number of days to flowering to 126 and 121 days, in the 1<sup>st</sup> and 2<sup>nd</sup> seasons, respectively, as compared with the non-saline treatment (control) (95 and 90 days, in the same order), while, it decreased the number of flowers/plant to 3 flowers/plant and an inflorescence diameter of 2.75 cm, in both seasons. Meanwhile, the number of flowers/plant and inflorescence diameter significantly increased in non-saline (control) to 9 flowers/plant and 4.15 cm, respectively, in both seasons. These results were similar to those of Nofal *et al.* (2015) and Shanan (2015), who

indicated that value of flowering characteristics significantly decreased in *Calendula officinalis* and *Matthiola incana* plants with increasing salt concentrations. This reduction in value of flowering characteristics may be due to the effect of salinity which inhibits photosynthesis of plants via changes of chlorophyll contents and components and damage of the photosynthetic apparatus. It also inhibits the photochemical activities and decreases the activities of enzymes (Mazhar *et al.*, 2012).

As shown in (Fig. 1) *L. indica* plants sprayed with the *Moringa* leaves extract recorded the significantly highest inflorescence number /plant (15 flowers/plant) and, inflorescence diameter (7.02 cm) and decreased the number of days to flowering (77.5 days) in both seasons, followed by plants sprayed with seaweed extract as compared with humic acid and Hoagland nutrient solution or control treatments under non-saline (control) and also, in saline conditions which the highest sea salt concentration (9.37 dS/m) recorded highest significant inflorescence number / plant (8.50 flowers/plant) and, inflorescence diameter (5.08 cm) and decreased the number of days to flowering (92.5 days), in both seasons. This supports the report of Culver *et al.* (2012) and Bashir *et al.* (2014) who reported that, application of the *Moringa* leaf extract after planting significantly increased number of flowers of the tomato plants as a result of its rich contents of different nutrients and, plant hormone like auxin and cytokinin.

#### Chemical composition

**Photosynthetic pigments:** Total chlorophyll and carotenoid contents under control condition were higher as compared to sea salt stressed treatments (Table 6). Total chlorophyll and carotenoid contents decreased when plants were subjected to sea salt stress. However, the maximum reduction in total




Fig. 1. Floral characters [number of days to flowering (a), inflorescence number /plant (b) and inflorescence diameter (c) of *L. indica* as affected by natural exogenous foliar applications under non-saline and sea salt concentrations during two seasons, 2014 (1<sup>st</sup>) and 2015 (2<sup>nd</sup>). Columns are significantly different at  $P = 0.05$  vertical bars represent  $\pm$ SE. S0=control, S1= 3.12 dS/m, S2= 6.25 dS/m, S3= 9.37 dS/m. C= distilled water as control, HNS= Hoagland nutrient solution, MLE= *Moringa* leaves extract, SE= seaweed, HA= humic acid.

chlorophyll and carotenoid contents were observed at 9.37 dS/m. That reduction of photosynthesis under sea salt stress could be attributed to the formation of proteolytic enzymes such as chlorophyllase, destruction of chlorophyll molecules by ROS, decline in membrane permeability, reducing water availability and nutrients particularly magnesium (Sharma and Hall, 1992; Kalaji *et al.*, 2011). Such inhibition in the photosynthesis after irrigation with sea water is also reported by Soliman *et al.* (2015) in *Moringa peregrine*.

Foliar spray of MLE led to the highest increment of total chlorophyll and carotenoid contents (2.56 and 1.29 mg/g f.w., in both seasons) in untreated plants (Table 6). However, total chlorophyll and carotenoid contents of crape myrtle plants significantly decreased with sea salt treatment at 9.37 dS/m and these increments were 1.70 and 0.86 mg/g f.w., in both seasons, as compared to non-saline (control) plants (0.79 and 0.40 mg/g f.w., in both seasons). The foliar spray MLE partially alleviates this reduction and minimized the reduction in chlorophyll contents. In addition, MLE maintained the chlorophyll in higher concentrations under salinity due to presence of zeatin-like cytokinin in MLE which prevents premature leaf senescence, induced cytokinin biosynthesis, and maintains higher leaf area for photosynthetic activity. It also contained calcium four times more than milk, potassium three times of banana and ascorbic acid seven times of orange and that led to reduce concentration of Na<sup>+</sup> ions and increased concentrations of K<sup>+</sup> and Ca<sup>2+</sup> ions coupled with the increased concentrations of carotenoids (Rady *et al.*, 2013; Zaki and Rady, 2015). These results corroborate findings of Azra *et al.* (2013).

#### Total carbohydrates, proline content and total soluble

Table 6. Chemical compositions of *L. indica* affected by exogenous foliar applications under different sea salt concentrations during two seasons, 2014 (1<sup>st</sup>) and 2015 (2<sup>nd</sup>)

Treatments		Total chlorophyll (mg/g f.w.)		Carotenoids (mg/g f.w.)		Total carbohydrates (%)		Proline content (μ moles /g f.w.)		Total phenol content (mg/g f.w.)	
		1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>
S0	C	1.48	1.50	0.75	0.77	15.00	17.67	17.00	19.33	0.52	0.55
	HNS	2.29	2.33	1.15	1.17	19.33	20.00	21.67	23.67	0.58	0.64
	MLE	2.54	2.58	1.28	1.29	25.00	28.00	26.33	28.00	0.64	0.70
	SE	2.42	2.49	1.21	1.26	22.67	24.67	25.67	27.00	0.59	0.69
	HA	2.29	2.34	1.14	1.18	20.33	22.00	24.00	26.00	0.56	0.62
S1	C	1.26	1.28	0.63	0.64	16.67	18.67	18.00	20.00	0.54	0.59
	HNS	2.11	2.16	1.06	1.08	20.00	21.00	22.33	25.00	0.58	0.66
	MLE	2.29	2.37	1.16	1.19	27.00	28.33	28.00	30.33	0.67	0.72
	SE	2.20	2.22	1.11	1.11	25.00	25.67	26.67	27.67	0.64	0.70
	HA	2.16	2.20	1.08	1.11	22.67	24.67	25.33	27.00	0.61	0.68
S2	C	1.16	1.19	0.58	0.60	21.33	25.00	21.67	24.67	0.59	0.63
	HNS	1.54	1.75	0.77	0.88	23.33	25.33	26.67	31.00	0.64	0.69
	MLE	1.96	2.18	0.99	1.08	31.67	34.00	34.67	37.33	0.72	0.78
	SE	1.83	2.10	0.92	1.05	28.33	31.33	34.00	36.00	0.71	0.75
	HA	1.70	1.91	0.86	0.96	27.33	29.00	31.33	35.00	0.65	0.72
S3	C	0.76	0.81	0.38	0.41	25.33	28.00	24.33	26.67	0.63	0.66
	HNS	1.22	1.30	0.61	0.65	28.33	32.33	31.00	32.67	0.69	0.72
	MLE	1.63	1.77	0.82	0.89	35.67	38.67	41.67	45.33	0.79	0.84
	SE	1.56	1.62	0.78	0.81	33.00	36.67	39.67	42.67	0.74	0.77
	HA	1.35	1.54	0.68	0.77	30.00	35.00	38.33	39.67	0.71	0.73
LSD (P=0.05)		0.06	0.06	0.03	0.03	1.71	1.16	1.90	1.83	0.03	0.04
		0.05	0.04	0.03	0.03	0.88	1.10	1.35	1.15	0.02	0.03
		0.11	0.09	0.05	0.06	1.77	2.20	2.70	2.30	0.05	0.07

S0=control, S1= 3.12 dS/m, S2= 6.25 dS/m, S3= 9.37 dS/m. C= distilled water as control, HNS= Hoagland nutrient solution, MLE= *Moringa* leaves extract, SE= seaweed, HA= humic acid.

**phenols:** It is obvious that total carbohydrates, proline content and total soluble phenols significantly increased with increasing levels of applied sea salt treatments (Table 6). Total carbohydrates, proline content and total soluble phenols in leaves of *L. indica* plants reached its maximum, in both seasons, in the plants irrigated by 9.37 dS/m sea salt. Many plants under salt stress accumulated starch and soluble carbohydrates as a response to impaired carbohydrate utilization, low cell division and growth since carbohydrate is a source of energy and osmolyte so it might play an important role in increasing the internal osmotic pressure (Zidan and Al-Zahrani, 1994; Miandoab *et al.*, 2015). Also, in response to salinity stress a large number of plant species accumulate proline, which may play a role in defense against salinity stress, cell osmotic adjustment, membrane stabilization and detoxification of injurious ions proline was considered as an indicator for some mechanism to resist the salinity stress (Ashraf and Foolad, 2007; Jampeetong and Brix, 2009). In addition, plants release phenolic metabolite which is induced in response to salinity and may strongly affect particularly leaf polyphenols and act as antioxidants to protect the plant against oxidative stress (Keutgen and Pawelzik, 2009; Radi *et al.*, 2013). These results are in harmony with many researches which concluded that the higher level of carbohydrate are associated with proline and total phenol (Deepika *et al.*, 2015; Mohsen *et al.*, 2015; Al-Hassan *et al.*, 2015) and might be a good strategy to increase salt tolerance.

Total carbohydrates, proline content and total soluble phenols significantly increased in salt-stressed crape myrtle plants by the exogenous applications of MLE (Table 6). This may be attributed to that MLE is excellent source for minerals, soluble sugars, amino acids, some antioxidants and a high cytokinin (zeatin & zeatin riboside) which causes increases in the concentrations of

Table 7. Macronutrients of *L. indica* affected by exogenous foliar applications under different sea salt concentrations during two seasons, 2014 (1<sup>st</sup>) and 2015 (2<sup>nd</sup>)

Treatments		N%		P%		K%		Na%		Ca%	
		1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>
S0	C	1.89	1.94	0.21	0.26	1.51	1.56	0.44	0.41	0.47	0.53
	HNS	2.29	2.36	0.23	0.29	1.57	1.59	0.38	0.37	0.53	0.58
	MLE	2.44	2.53	0.30	0.34	1.67	1.70	0.29	0.26	0.60	0.67
	SE	2.36	2.48	0.29	0.30	1.63	1.68	0.36	0.32	0.56	0.64
	HA	2.30	2.41	0.26	0.29	1.60	1.63	0.37	0.35	0.54	0.60
S1	C	1.82	1.85	0.17	0.22	1.48	1.54	0.52	0.47	0.70	0.78
	HNS	2.24	2.30	0.21	0.25	1.53	1.59	0.47	0.44	0.88	0.97
	MLE	2.41	2.45	0.28	0.31	1.64	1.68	0.39	0.35	0.99	1.17
	SE	2.31	2.41	0.27	0.28	1.62	1.65	0.41	0.39	0.96	1.08
	HA	2.28	2.32	0.22	0.26	1.58	1.61	0.44	0.43	0.92	1.01
S2	C	1.56	1.61	0.11	0.18	1.37	1.43	0.82	0.88	1.22	1.29
	HNS	1.60	1.72	0.16	0.19	1.46	1.51	0.80	0.76	1.51	1.57
	MLE	2.07	2.17	0.21	0.26	1.58	1.63	0.64	0.59	1.55	1.79
	SE	1.97	2.01	0.19	0.22	1.52	1.62	0.68	0.62	1.61	1.71
	HA	1.83	1.96	0.18	0.19	1.47	1.54	0.74	0.67	1.57	1.66
S3	C	1.19	1.23	0.07	0.11	1.25	1.30	1.13	1.06	1.54	1.63
	HNS	1.55	1.62	0.11	0.15	1.37	1.40	1.07	1.02	1.71	1.88
	MLE	1.94	2.05	0.17	0.20	1.42	1.50	0.96	0.90	1.95	2.03
	SE	1.87	1.95	0.15	0.19	1.44	1.47	0.99	0.96	1.91	3.00
	HA	1.79	1.84	0.13	0.16	1.40	1.45	1.04	1.00	1.55	1.90
LSD ( $P=0.05$ )		0.07	0.02	0.04	0.03	0.03	0.06	0.09	0.02	0.14	0.24
		0.05	0.04	0.03	0.02	0.05	0.05	0.04	0.06	0.10	0.35
		0.11	0.09	0.07	0.05	0.10	0.10	0.12	0.09	0.21	0.70

S0=control, S1= 3.12 dS/m, S2= 6.25 dS/m, S3= 9.37 dS/m. C= distilled water as control, HNS= Hoagland nutrient solution, MLE= *Moringa* leaves extract, SE= seaweed, HA= humic acid.

proline and total soluble phenols, in turn, protect plants against the ROS generation, osmotic adjustment and membrane injury, or may result in the synthesis of other substances having a protective effect on plants grown under salt stress (Thakur and Sharma, 2005; Rady *et al.*, 2015). In addition, Munoz *et al.* (2008) concluded that zeatin riboside affected mainly the mobilization of carbohydrates as well as distributed to the sink where more carbohydrates are needed to cater the needs of rapidly increasing growth. The protective role of *Moringa* leaf extract against ROS that are formed during salt stress was reported by Iqbal (2014).

**Macro-nutrients:** Sea salt stress had a significant effect on all macro-nutrients; N, P, K, Ca and, Na%; in both seasons. Shoot N (1.21), P (0.24) and K (1.54) % decreased significantly ( $P \leq 0.01$ ) as sea salt increased, in both seasons, than those of the control (1.92, 0.09 and 1.28%, in both seasons), while Na and Ca% increased with increasing sea salt concentrations. Maximum Na and Ca% was recorded at 9.37 dS/m, 1.10 and 1.59% in both seasons, as compared to control (0.43 and 0.50 %, in both seasons) (Table 7). Sea salt concentrations causing conflict in nutrient homeostasis by competing with the uptake of other essential nutrients. Also, the excessive Na<sup>+</sup> decreases the uptake of other essential nutrients such as N, P, K<sup>+</sup> and Ca<sup>++</sup> (Shahbaz *et al.*, 2013; Parihar *et al.*, 2014). These results are in harmony with those of Cornacchione and Suarez (2015) and Ferreira *et al.* (2015).

Application of MLE decreased the concentration of Na<sup>+</sup> and increased concentration of N, P, K<sup>+</sup> and, Ca in leaves against control, under normal as well as sea salt concentrations (Table 7). This result could be due to the important role of *Moringa* leaf extract which contain proteins, amino acids and various phenolics and zeatin, ascorbic acid, phenolic compounds, sugars, and minerals (*i.e.* calcium, magnesium, sodium, iron, phosphorus and

potassium); hence, it is used as a natural plant growth enhancer (Anwar *et al.*, 2007). Therefore, it has the ability to reduce the adverse effect of salinity and to protect the plant against the build-up of toxic ions by maintenance of the ionic homeostasis under salt stress and this makes N, P, K, Ca more available to the plant. These results are parallel with the finding of Abd El-Hamied and El-Amry (2015).

**Enzyme activity:** The activity of CAT, SOD and APX increased significantly with increasing sea salt concentrations (Fig. 2). Untreated plants (control) showed the lowest significant CAT (1.42), SOD (3.91) and, APX (2.12 units/min/mg protein), in both seasons. In plants under salt stress activities of the antioxidative enzymes (CAT, SOD and, APX) increase to avoid the damage caused by oxidative stress that induced by the elevated levels of reactive oxygen species (ROS) such as hydrogen peroxide (Sharma *et al.*, 2012). SOD constitutes the first line of defense against ROS by reducing the O<sub>2</sub><sup>-</sup> radical to H<sub>2</sub>O<sub>2</sub> (Alscher *et al.*, 2002). In addition, APX is considered as one of the most important enzymes in the reduction of this reactive molecule (Feierabend, 2005). This was in conformity with the earlier findings of Muthulakshmi *et al.* (2015).

*Moringa* leaves extract had the superiority in increasing enzyme activities (CAT, SOD and APX) in sea salt stressed or unstressed plants, followed by seaweed extract, humic acid, Hoagland nutrient solution which was more than the control. *Moringa* leaves extract increased enzyme activities at 9.37 dS/m by 3.14, 14.69, and 4.49, respectively, in both seasons, as compared with untreated plants under the same level of sea salt stress (1.89, 7.48, and 2.81 units/min/mg protein, respectively, in both seasons). This increase may be attributed to that *Moringa* leaves extract which contains several photochemicals, flavonoid pigments such as




Fig. 2. Enzyme activity [CAT (a), SOD (b) and APX (c)] of *L. indica* as affected by natural exogenous foliar applications under non-saline and sea salt concentrations during two seasons, 2014 (1<sup>st</sup>) and 2015 (2<sup>nd</sup>) Columns are significantly different at  $P=0.05$  vertical bars represent  $\pm$ SE.

S0=control, S1= 3.12 dS/m, S2= 6.25 dS/m, S3= 9.37 dS/m. C= distilled water as control, HNS= Hoagland nutrient solution, MLE= *Moringa* leaves extract, SE= seaweed, HA= humic acid.



kaempferol, rhamnetin, isoquercitrin and kaempferitrin and, rich in zeatin and a group of the glycoside compounds, glucosinolates and isothiocyanates, which may enhance the formation of the photochemical compounds and that led to increase in the antioxidant activity. Moreover, to control the ROS production, repair and mitigate damage initiated by reactive oxygen, plants develop a complex antioxidant system enabling survival under abiotic stress, so MLE is proved to be the most effective PGR in reducing plant exposure to certain stress (Rajanandh *et al.*, 2012). Several studies have indicated such results (Azra *et al.*, 2013; Aslam *et al.*, 2015).

This study indicated that *Moringa* leaves extract (MLE) applied to *L. indica* plants at different concentrations of sea salt improved vegetative growth, floral characters and chemical compositions which enhanced the tolerance of sea salt stress by increasing the proline, total phenolic contents and activation of the antioxidant enzyme system of SOD, CAT and APX. *Moringa* leaves extract can be used as a possible supplement or substitute to inorganic or chemical fertilizer as growth enhancer and under different sea salt concentrations. Therefore, MLE can be recommend to be used commercially as environment friendly and economically cheap fertilizer especially in arid and semi-arid regions.

## Reference

- Abd El-Hamied, S.A. and E.I. El-Amary, 2015. Improving growth and productivity of "Pear" trees using some natural plants extracts under North Sinai conditions. *IOSR Journal Agriculture Veterinary Science*, 8(1): 1-9.
- Afzal, I., B. Hussain, S.M.A. Basra and H. Rehman, 2012. Priming with MLE reduces imbibitional chilling injury in spring maize. *Seed Science Technology*, 40(2): 271-276.
- AL Hassan, M., M.M. Fuertes, J.R. Sánchezf, O. Vicente and M. Boscaiu, 2015. Effects of salt and water stress on plant growth and on accumulation of osmolytes and antioxidant compounds in cherry tomato. *Not. Bot. Hort. Agrob.*, 43(1): 1-11.
- AL Khayri, J.M. 2002. Growth, proline accumulation and ion content in sodium chloride stressed callus of date palm. *In Vitro Cellular Developmental Biology*, 38: 79-82.
- Allen, S.F., H.F. Grimshaw and A.B. Rowl, 1984. Chemical analysis. In: *Methods in Plant Ecology*. P.D. Moor and S.B. Chapman (eds.). Blackwell, Oxford. p. 185-344.
- Alscher, R.G., N. Erturk and L.S. Heath, 2002. Role of superoxide dismutases (SODs) in controlling oxidative stress in plants. *J. Experimental Biology*, 53: 1331-1341.
- Anderson, N.O. 2007. *Flower Breeding and Genetics*. Springer, Dordrecht, pp. 439-457.
- Anonymous, 1962. *The Wealth of India, Publication and Information Directorate*. Council of Scientific and Industrial Research, New Delhi, India, 7: 24.
- Anwar F., S. Latif, M. Ashraf and A.H. Gilani, 2007. *Moringa oleifera*: A food plant with multiple medicinal uses. *Phytother. Res.*, 21: 17-25.
- Ashraf, M. and M.R. Foolad, 2007. Roles of glycinebetaine and proline in improving plant abiotic stress tolerance. *Environ. Expt. Bot.*, 59: 206-216.
- Aslam, M., B. Sultana, F. Anwar and H. Munir, 2015. Foliar spray of selected plant growth regulators affected the biochemical and antioxidant attributes of spinach in a field experiment. *Turkish J. Agriculture Forestry*, 39: 1-10.
- Azra, Y., S.M.A. Basra, R. Ahmad and A. Wahid, 2012. Performance of late sown wheat in response to foliar application of *Moringa oleifera* Lam. leaf extract. *Chil. J. Agric. Res.*, 72: 92-97.
- Azra, Y., S.M.A. Basra, F. Muhammad, ur R. Hafeez, H. Nazim and R.A. Habib, 2013. Exogenous application of *Moringa* leaf extract modulates the antioxidant enzyme system to improve wheat performance under saline conditions. *Plant Growth Regul.*, 69: 225-233.
- Bashir, K.A., J.A. Bawa and I. Mohammed, 2014. Efficacy of leaf extract of drumstick tree (*Moringa oleifera* Lam.) on the growth of local tomato (*Lycopersicon esculentum*). *J. Pharmacy and Biological Sciences*, 9(4): 74-79.
- Bates, L.S., R.P. Waldern and L.D. Teare, 1973. Rapid determination of free proline under water stress studies. *Plant and Soil*, 39: 205-207.
- Battacharya, D., M.Z. Babgohari, P. Rathor and B. Prithiviraj, 2015. Seaweed extracts as biostimulants in horticulture. *Scientia Hort.*, 196: 39-48.
- Cabrera, R.I. and D.R. Devereaux, 1999. Crapemyrtle post-transplant growth as affected by nitrogen nutrition during nursery production. *Journal American Society Horticultural Science*, 124: 94-98.
- Chance, B. and A.C. Maely, 1955. Assay of catalase and peroxidase methods. *Enzymology*, 2: 755-784.
- Chapman, H.D. and P.F. Pratt, 1961. *Methods of Analysis for Soil, Plants and Water*. University of California, Division of Agricultural Science, Berkeley, CA, USA, p. 56-63.
- Chopra, R.N., S.C. Nagar and I.C. Chopra, 1956. In *Glossary of Indian Medicinal Plants*. Council of Scientific and Industrial Research, New Delhi. 1:197.
- Conti, E., A. Litt, P.G. Wilson, S.A. Graham, B.G. Briggs, L.A.S. Johnson and K.J. Sytsma, 1997. Interfamilial relationships in Myrtales: molecular phylogeny and patterns of morphological evolution. *Syst. Bot.*, 22: 629-647.
- Cornacchione, M.V. and D.L. Suarez, 2015. Emergence, forage production and ion relations of alfalfa in response to saline waters. *Crop Sci.*, 55: 444-457.
- Culver, M., T. Fanuel and A.Z. Chiteka, 2012. Effect of *Moringa* extract on growth and yield of tomato. *Greener Journal Agricultural Sciences*, 2(5): 207-211.
- Deepika, U.K.V., P.S. Manogya and S.S. Preeti, 2015. Influence of salinity on morpho-physiological characters of german chamomile (*Matricaria chamomilia* L.). *World Journal Pharmacy Pharmaceutical Sciences*, 4(10): 762-775.
- Diab, Y., K. Atalla and K. Elbanna, 2012. Antimicrobial screening of some Egyptian plants and active flavones from *Lagerstroemia indica* leaves. *Drug. Dis. Therap.*, 6(Suppl 4): 212-217.
- DuBois, M., F. Smith, K.A. Gilles, J.K. Hamilton and P.A. Rebers, 1956. Colorimetric method for determination of sugars and related substances. *Annal. Chem.*, 28(3): 350 -356.
- Feierabend, J. 2005. Catalases in plants: Molecular and functional properties and role in stress defense. In: *Antioxidants and Reactives Oxygen Species in Plants*. N. Smirnov (ed.), Blackwell Publishing, London, UK. p. 101-140.
- Ferreira, J.F.S., M.V. Cornacchione, X. Liu and D.L. Suarez, 2015. Nutrient composition, forage parameters and antioxidant capacity of alfalfa (*Medicago sativa*, L.) in response to saline irrigation water. *Agriculture*, 5: 577-597.
- Foxcroft, L.C., D.M. Richardson and J.R.U. Wilson, 2007. Ornamental plants as invasive aliens: problems and solutions in Kruger National Park. *South Africa. Environmental Management*, 41(1): 32-51.
- Gamble, J.S. 1972. *A Manual of Indian Timbers*. 2<sup>nd</sup> ed. Bishen Singh, Mahendra Pal Singh, Dehradun, India, p 868.
- Giannopolitis, N.C. and S.K. Ries, 1977. Superoxide dismutases: I. Occurrence in higher plants. *Plant Physiology*, 68: 548-552.
- Hoagland, D.R. and D.I. Arnon, 1950. *The Water-Culture Method for Growing Plants Without Soil*. Circular 347. Agricultural Experiment Station, University of California, Berkeley, 1-32.
- Howladar, S.M. 2014. A novel *Moringa oleifera* leaf extract can mitigate the stress effects of salinity and cadmium in bean (*Phaseolus vulgaris* L.) plants. *Ecotoxicol. Environ. Saf.*, 100: 69-75.

- Iqbal, M.A. 2014. Role of *Moringa*, *Brassica* and *Sorghum* water extracts in increasing crops growth and yield: A Review. *American-Eurasian J. Agric. Environ. Sci.*, 14(11): 1150-1158.
- Jackson, M.L. 1967. *Soil Chemical Analysis*. Prentice-Hall, India. p. 144-197.
- Jampeetong, A. and H. Brix, 2009. Nitrogen nutrition of *Salvinia natans*: Effects of inorganic nitrogen form on growth, morphology, nitrate reductase activity and uptake kinetics of ammonium and nitrate. *Aquat. Bot.*, 90: 67-73.
- Kalaji, H.M., B.K. Govindjee, J. Koscielniak and K. Zuk-Goaszewska, 2011. Effects of salt stress on photosystem II efficiency and CO<sub>2</sub> assimilation of two Syrian barley landraces. *Environ Exp. Bot.*, 73: 64-72.
- Keutgen, A.J. and E. Pawelzik, 2009. Impacts of NaCl stress on plant growth and mineral nutrient assimilation in two cultivars of strawberry. *Environ. Experi. Bot.*, 65(2-3): 170-176.
- Kotnala, M., G.S. Chakraborty and A. Mazumder, 2013. *Lagerstroemia* Species: A Current Review. *International Journal PharmTech Research*, 5(3): 906-909.
- Little, T.M. and F.J. Hills, 1978. *Agricultural Experimentation Design and Analysis*. John Wiley & Sons, Inc., New York, USA: pp. 53-63.
- Masharani, U. and M.S. German, 2011. Pancreatic hormones and diabetes mellitus. In: *Greenspan's Basic and Clinical Endocrinology*. D.G. Gardner and D. Shoback, (eds.), 9<sup>th</sup> ed., McGraw-Hill, New York, United States. p. 573-655.
- Mazhar, A.M., S.I. Shedeed, G. Nahed, N.G. Abdel-Aziz and M.H. Mahgoub, 2012. Growth, flowering and chemical constituents of *Chrysanthemum indicum* L. plant in response to different levels of humic acid and salinity. *J. Applied Sciences Research*, 8(7): 3697-3706.
- Miandoab, L.Z., M.B. Bagherieh-Najjar, M.A. Hejazi and N. Chaparzadeh, 2015. The psy transcript level and cell composition of *Dunaliella salina* under salinity. *International Journal Biosciences*. 6(7): 34-43.
- Mohsen, A.A., M.A. Elhaak, E.A.M. Hamada and F.E. El-Gebaly, 2015. Adaptation potential of two common halophytes to salinity stress in the salt marshes of lake Burullus in Egypt. *International J. Advances Pharmacy, Biology Chemistry*, 4(4): 809-820.
- Moore, L.M. and F.D. Walker, 2014. Plant Guide: Crape myrtle *Lagerstromia indica*. USA: USDA Natural Resources Conservation Service. 1-4, [http://plants.usda.gov/plantguide/pdf/pg\\_lain.pdf](http://plants.usda.gov/plantguide/pdf/pg_lain.pdf)
- Moussa, H.R. and M.A.-E. Hassan, 2016. Growth enhancers to mitigate salinity stress in *Vicia faba*. *International Journal of Vegetable Science*, 22(3): 243-250.
- Moyo, B., P.J. Masika, A. Hugo and V. Muchenje, 2011. Nutritional characterization of *Moringa (Moringa oleifera* Lam) leaves. *African Journal Biotechnology*, 10(60): 12925 -12933.
- Muthulakshmi, S.M., S.G. Gurulakshmi and S. Rajathi, 2015. Effect of salt stress on physiological and biochemical characteristics in *Solanum nigrum* L. *International J. Science and Research (IJSR)*, 4(3): 567-571.
- Munns, R., R.A. James and A. Lauchli, 2006. Approaches to increasing the salt tolerance of wheat and other cereals. *J. Exp. Bot.*, 57: 1025-1043.
- Moran, R. 1982. Formula for determination of chlorophyll pigments extracted with N.N. dimethyl formamide. *Plant Physiol.*, 69: 1371-1381.
- Munoz, J.L., L. Martin, G. Nicolas and N. Villalobos, 2008. Influence of endogenous cytokinins on reverse mobilization in cotyledons of *Cicer arietinum* L. *J. Agric. Res.*, 93(3): 1011-1016.
- Nofal, F.H., M.U. El-Segai and E.A. Seleem, 2015. Response of *Calendula officinalis* L. Plants to growth stimulants under salinity stress. *American-Eurasian J. Agric. Environ. Sci.*, 15(9): 1767-1778.
- Orwa, C., A. Mutua, R. Kindt, R. Jamnadass and A. Simons, 2009. *Agroforestry Database: a tree reference and selection guide* (version 4.0), World Agroforestry Centre, Kenya. (<http://www.worldagroforestry.org/af/treedb/>)
- Oviedo, P.R., O.P. Herrera and M.G. Caluff, 2012. National list of invasive and potentially invasive plants in the Republic of Cuba-2011. (Lista nacional de especies de plantas invasoras y potencialmente invasoras en la República de Cuba-2011). *Bissea: Boletín sobre Conservación de Plantas del Jardín Botánico Nacional de Cuba*, 6 (Special Issue 1): 22-96.
- Parihar, P., S. Singh, R. Singh, V.P. Singh and S.M. Prasad, 2014. Effect of salinity stress on plants and its tolerance strategies: a review. *Environ Sci Pollut Res Int.*, 22(6): 4056-4075.
- Piper, C.S. 1950. *Soil and Plant Analysis*. Univ. Adelied. Interscience Published, Inc. New York, pp: 258-275.
- Price, L.M. 2007. *The Moringa tree*. ECHO Technical Note. Online available at [http://www.echotech.org/mambo/index.php?option=com\\_docman&task=doc\\_view&gid=170:327-336](http://www.echotech.org/mambo/index.php?option=com_docman&task=doc_view&gid=170:327-336).
- Pregl, P. 1945. *Quantitative Organic Microanalysis*. Appl. Churchill Publishing Co., London, 4<sup>th</sup> Ed., p. 78-82.
- Radi, A.A., F.A. Farghaly and A.M. Hamada, 2013. Physiological and biochemical responses of salt-tolerant and salt-sensitive wheat and bean cultivars to salinity. *J. Biol. Earth. Sci.*, 3(1): 72-88.
- Rady, M.M., V.C. Bhavya and S.M. Howladar, 2013. Common bean (*Phaseolus vulgaris* L.) seedlings overcome NaCl stress as a result of presoaking in *Moringa oleifera* leaf extract. *Scientia Horticulturae*, 162: 63-70.
- Rady, M.M. and G.F. Mohamed, 2015. Modulation of salt stress effects on the growth, physio-chemical attributes and yields of *Phaseolus vulgaris* L. plants by the combined application of salicylic acid and *Moringa oleifera* leaf extract. *Scientia Horticulturae*, 193: 105-113.
- Rady, M.M., G.F. Mohamed, A.M.A. Bdalla and Y.H.M. Ahmed, 2015. Integrated application of salicylic acid and *Moringa oleifera* leaf extract alleviates the salt-induced adverse effects in common bean plants. *Journal of Agricultural Technology*, 11(7):1595-1614.
- Rady, M.M., T.A. Abd El-Mageed, H.A. Abdurrahman and A.H. Mahdi, 2016. Humic acid application improves field performance of cotton (*Gossypium barbadense* L.) under saline conditions. *The Journal of Animal Plant Sciences*, 26(2): 487-493.
- Rajanandh, M.G., M.N. Satishkumar, K. Elango and B. Suresh, 2012. *Moringa oleifera* Lam. A herbal medicine for hyperlipidemia: a pre-clinical report. *Asian Pac. J. Trop. Dis.*, 2: S790-S795.
- Randall, R.P. 2012. *A Global Compendium of Weeds*. 2<sup>nd</sup> Edition. Department of Agriculture and Food, Western Australia, 571.
- Rao, M.V., G. Paliyath and D.P. Ormrod, 1996. Ultraviolet-B radiation and ozone-induced biochemical changes in the antioxidant enzymes of *Arabidopsis thaliana*. *Plant Physiology*, 110: 125-136.
- Rehman, H., M.Q. Nawaz, S.M.A. Basra, I. Afzal, Y. Azra and F.U., Hassan, 2014. Seed priming influence on early crop growth, phenological development and yield performance of linola (*Linum usitatissimum* L.). *Journal Integrative Agriculture*, 13(5): 990-996.
- Sadak, M.Sh. and M.G. Dawood, 2014. Role of ascorbic acid and  $\alpha$  tocopherol in alleviating salinity stress on flax plant (*Linum usitatissimum* L.). *Journal Stress Physiology Biochemistry*, 10(1): 93-111.
- Shahbaz, M., Z. Mushtaq, F. Andaz and A. Masood, 2013. Does proline application ameliorate adverse effects of salt stress on growth, ions and photosynthetic ability of eggplant (*Solanum melongena* L.). *Sci. Hort.*, 164: 507-511.
- Shanan, N.T. 2015. Alleviation of salt stress by stimulative compounds in *Matthiola incana* L. plants. *International J. Advanced Research*, 3(9): 665-675.
- Sharma, P.K. and D.O. Hall, 1992. Changes in carotenoid composition and photosynthesis in sorghum under high light and salt stress. *J. Plant Physiol.*, 140: 661-666.
- Sharma, P., A.B. Jha, R.S. Dubey and M. Pessarakli, 2012. Reactive oxygen species, oxidative damage and antioxidative defense mechanism in plants under stressful conditions. *Journal Botany*, vol. 2012, Article ID 217037, 26 pages.

- Soliman, A.Sh., S.A. El-Feky and E. Darwish, 2015. Alleviation of salt stress on *Moringa peregrina* using foliar application of nanofertilizers. *Journal Horticulture Forestry*, 7(2): 36-47.
- Swain, T. and W.E. Hillis, 1959. Phenolic constituents of *Prunus domestica* I quantitative analysis of phenolic constituents. *J. Science Food Agriculture*, 10: 63-68.
- Tester, M. and R.J. Davenport, 2003. Na<sup>+</sup> tolerance and Na<sup>+</sup> transport in higher plants. *Ann. Bot.*, 91: 503-527.
- Thakur, M. and A.D. Sharma, 2005. Salt-stress-induced proline accumulation in germinating embryos: evidence suggesting a role of proline in seed germination. *Journal Arid Environments*, 62: 517-523.
- Tobe, H. and P.H. Raven, 1990. Comparative reproductive morphology of *Orias* and *Lagerstroemia* (Lythraceae). *Flora*, 184: 177-185.
- Vitoria, A.P., P.J. Lea and R.A. Azevedo, 2001. Antioxidant enzymes responses to cadmium in radish tissues. *Phytochemistry*, 57: 701-710.
- Yang, E.J., J.S. Lee, B.B. Song, C.Y. Yun, D.H. Kim and I.S., Kim, 2011. Anti-inflammatory effects of ethanolic extract from *Lagerstroemia indica* on airway inflammation in mice. *Journal of Ethnopharmacology*, 136(3): 422-427.
- Yeung, L.T. 2010. Biology Project, *Introduction of Hong Kong Plants*, 6S, 13: 13-14.
- Zaki, S.S. and M.M. Rady, 2015. *Moringa oleifera* leaf extract improves growth, physiochemical attributes, antioxidant defence system and yields of salt-stressed *Phaseolus vulgaris* L. plants. *International J. Chem. Tech. Research*, 8(11): 120-134.
- Zidan, M.A. and H.S. Al-Zahrani, 1994. Effect of NaCl on the germination seedling and some metabolic changes in Sweet Basil (*Ocimum basilicum*). *Pak. J. Sci. Ind. Res.*, 37: 541-543.

---

Received: November, 2016; Revised: January, 2017;

Accepted: February, 2017